

Forest Herbicide Effectiveness on Mississippi Tree and Shrub Species

Regarding forest herbicide efficacy, one of the most common concerns is how well an herbicide controls a specific tree or shrub species. Obviously, if you purchase herbicide and spend the time required to apply it, you want effective control of the targeted stems. While individual herbicide labels contain some species-specific information, the list of woody species and their tolerance to that particular herbicide is typically limited.

The table on the next page provides a list of commonly used forest herbicides and their associated effectiveness in controlling certain species. Susceptibility level is shown for each herbicide and each woody species specifically. These levels of susceptibility were established using the references below, herbicide labels, and personal field research experience. These ratings are not absolute, as herbicide efficacy depends on multiple factors. However, these assessments provide expected effectiveness of listed herbicides under normal application conditions and proper application procedures.

Control is categorized as susceptible (S), intermediate (I), or tolerant (T). A susceptible ranking means that at least 80 percent control of species in that particular genus is expected. A ranking of intermediate means that 40–80 percent control can be expected. If you use an intermediate-ranked herbicide, an additional herbicide application will likely be needed. A ranking of tolerant means that herbicide will not provide control and should not be considered. If there is no ranking for a particular herbicide under a species grouping, there is not enough information available to offer an assessment.

References

- Cantrell, R. L. *A Guide to Silvicultural Herbicide Use in the Southern United States*. Auburn University, Auburn University School of Forestry. November 1985.
- Cunningham, K., Taylor, E., Barber, B., Holley, G., and Blazier, M. 2019. *Forestry Herbicide Prescriptions: Western Gulf Region*. University of Arkansas Cooperative Extension Service publication MP553.
- Heiligmann, R. B., and Krause, D. 2006. *Relative Effectiveness of Herbicides Commonly Used to Control Woody Vegetation in Forest Stands*. The Ohio State University Extension factsheet F-51-06.
- Miller, J. H., and Mitchell, R. J. 1990. *A Manual on Ground Applications of Forestry Herbicides*. USDA Forest Service. Management bulletin R8-MB21.

Effectiveness of forest herbicides by woody species group.

Herbicide	Maple	Alder	Serviceberry	Birch	Hornbeam	Hickory/Pecan	Hackberry/Sugarberry	Redbud	Dogwood	Hawthorn	Persimmon	Beech	Ash	Honeylocust	Walnut	Juniper	Sweetgum	Poplar	Osage orange	Mulberry	Blackgum	Hophornbeam	Sourwood	Pine	Sycamore	Cottonwood	Prunus species	Oak	Sumac	Black locust	Willow	Sassafras	Chinese tallow	Elm
Aminopyralid	S	S				S	S	S						S		S								I		S	S	S		S				S
Clethodim	T	T	T	T	T	T	T	T	T	T	T	T	T	T	T	T	T	T	T	T	T	T	T	T	T	T	T	T	T	T	T	T	T	T
Clopyralid	T	T	T	T	T	T	T	S	T	T	T	T	T	S	T		T	T			T	T	T	T	T	T	T		S		T		T	
Dicamba	I	S	S	S	S	S			S	S	T		S				T						I	S		S	I	I	S		S			S
Fosamine	T	S		S		T				T	S		S				S	T					T	S	S	S	I	S	S	S	T	T	S	I
Glyphosate	I	S	I	S	I	I	I	T	I	S	I	S	I	I	I		S	S	T	I	I	I	I	S	S	S	I	S	I	I	I	I	I	I
Hexazinone	I			S		I			I	S	T		I				S	T			T		T	T		S	I	S	I		S	T		S
Imazapyr	S	S	S	S	I	S	I	T	S	S	S	S	S	T	S		S	S	T	S	S	I	S	T ¹	S	S	S	S	S	T	S	S	S	T
Metsulfuron	S	S								S			S			S		I		S				T ²		S	S	S		S	S			S
Oxyflourfen	T	T	T	T	T	T	T	T	T	T	T	T	T	T	T	T	T	T	T	T	T	T	T	T	T	T	T	T	T	T	T	T	T	T
Picloram	S	S	S	S	S	S	S	S	I	S	S	S	I	S	S	S	I	S	S	S	S	S		I	S	S	S	S	S	S	S	S		I
Picloram + 2,4-D	S	S	S	S	S	S	S	S	S	S	S	S	S	S	S	S	I	S	S	S	S	S	S	I	S	S	S	S	S	S	S	S		S
Sulfometuron	T	T	T	T	T	T	T	T	T	T	T	T	T	T	T	T	T	T	T	T	T	T	T	T	T	S	T	T	I	T	T	T	T	T
Triclopyr (amine)	S	S		S	S	I	S	S	S	S	I	S	S	I	S	T	S	S	S	S	S	S	I	I	S	S	I	I	I	S	S	S	S	I
Triclopyr (ester)	I	S		S	S	I	S	S	S	I	I	S	I	I	S	I	S	S	S	S	S	S	I	I	S	S	I	I	I	S	S	S	S	S
2, 4-D	I	I		I	T	I	I		S	I	I	I	T	S	I				I	I	S	T				I	S	I	S	I	I	I	I	I

S = susceptible, I = intermediate, T = tolerant, no ranking = information too limited for susceptibility assessment.

¹Susceptible to Chopper Gen2.

²Will kill longleaf pine.

Publication 3415 (POD-04-26)

By A. Brady Self, PhD, Extension Professor, Forestry.



Copyright 2026 by Mississippi State University. All rights reserved. This publication may be copied and distributed without alteration for nonprofit educational purposes provided that credit is given to the Mississippi State University Extension Service.

Produced by Agricultural Communications.

Mississippi State University is an equal opportunity institution. Discrimination is prohibited in university employment, programs, or activities based on race, color, ethnicity, sex, pregnancy, religion, national origin, disability, age, sexual orientation, genetic information, status as a U.S. veteran, or any other status to the extent protected by applicable law. Questions about equal opportunity programs or compliance should be directed to the [Office of Civil Rights Compliance](#), 231 Famous Maroon Band Street, P.O. 6044, Mississippi State, MS 39762.

Extension Service of Mississippi State University, cooperating with U.S. Department of Agriculture. Published in furtherance of Acts of Congress, May 8 and June 30, 1914. ANGUS L. CATCHOT JR., Director